

| | | | |
|--|--|--|--|
| Surgeon | Dr.Gilmer | Procedure | Knee Arthroscopy Level 2 |
| Glove Size | 8 Bio, 7 ½ whites x2 | Position of Patient | Supine with Lateral Post |
| Skin Prep | Chloraprep Hibiclens wash and shaving and alcohol needs to be done in pacu. | | |
| Sutures Needles | 3-0 Nylon FS-1 | Drapes | Arthrotomy Pack, Extremity Sheet, Impervious U drapes x2, Impervious stockinette, Coban, 6" Esmark, Single Basin set |
| Dressings | Sm Xeroform, 4x4's, 4" Webril, ABD, Double 6" Ace | Instruments and Equipment | |
| | | Arthroscopy tray Arthroscope Athrex shaver Camera & Light Cord Pump and patient tubing 4.0 Dbl cut Cvd 18ga 1 ½" needle (for drawing Morphine) If meniscal repair: Arthrex knee scorpion (AR-12990 pulldown) On knee cart, hold: Stryker AIR (replaces S&N fast fix) on knee cart Inside/out repair: Sharpshooter Canulas (tray on knee cart) Sharpshooter handle (disposable, on knee cart) Hold: Torpedo Curved Knee Cart-have in room/extra supplies are on cart. | |
| Extra Items | | | |
| 0.5 % Marcaine with Epi @ beginning (10cc's usually), use same syringe for the following: Morphine 5 mg and 0.5% Ropivacaine plain 10 ml | | | |
| Notes | | | |
| 1. Foot Stool on Surgical side and shaver foot pedal on it. 2. If meniscus repair....no Epi 3. If repair, bring in knee cart 4. Uses straight hemostat after stab incision 5. 6 pictures per page, 1 copy Uses 15 blade in pack Room lights at ½ (Not out completely) Monitor placed at patient's opposite shoulder | | | |
| | | 7/11/17 KK | |